

# XMM-NEWTON SPECTRAL ANALYSIS OF SWIFT J1727.8-1613 ACROSS THE SOFT-TO-HARD TRANSITION.

Maïmouna Brigitte  
PhD student with Jiří Svoboda

**Collaborators:** Noel Castro Segura, María Díaz Trigo, Stefanie Fijma, Christian Knigge and Jakob Van den Eijnden



FACULTY  
OF MATHEMATICS  
AND PHYSICS  
Charles University



Astronomical  
Institute  
of the Czech Academy  
of Sciences

Credit John Paice

# TABLE OF CONTENTS

**01.**

## **THE SOURCE**

Swift J1727.8-1613

**02.**

## **OBSERVATIONS**

with XMM-Newton

**03.**

## **SPECTRAL MODELLING**


of the EPIC-pn and RGS  
spectra

**04.**

## **PROSPECTS WITH NEW ATHENA**

with X-IFU

## The source: the LMXB Swift J1727.8 – 1613



### ❏ Low Mass X-ray binary (LMXB)

- $d = 3.4 - 5.5^{+1.4}_{-1.1}$  kpc
- $i \leq 74^\circ$
- $P_{\text{orb}} = 10.8$  h

Burridge et al. 2025

Mata Sanchez et al. 2024, 2025

Wood et al. 2024


### Early K-type star

- $0.6 M_\odot < M_{\text{star}} < 0.9 M_\odot$

### Stellar Black hole

**Credits:** Hynes, <http://www.astro.soton.ac.uk/~rih/>.

# The source: the LMXB Swift J1727.8 – 1613



König et al. 2026

☐ Low Mass X-ray binary (LMXB)

- $d = 3.4 - 5.5^{+1.4}_{-1.1}\text{ kpc}$
- $i \leq 74^\circ$
- $P_{\text{orb}} = 10.8\text{ h}$

Burridge et al. 2025

Mata Sanchez et al. 2024, 2025


Wood et al. 2024

**Early K-type star**

- $0.6 M_{\odot} < M_{\text{star}} < 0.9 M_{\odot}$

**Stellar Black hole**

# The source: the LMXB Swift J1727.8 – 1613



radio jet

☐ Low Mass X-ray binary (LMXB)

- $d = 3.4 - 5.5^{+1.4}_{-1.1}$  kpc
- $i \leq 74^\circ$
- $P_{\text{orb}} = 10.8$  h

Burridge et al. 2025

Mata Sanchez et al. 2024, 2025

Wood et al. 2024

Early K-type star

- $0.6 M_\odot < M_{\text{star}} < 0.9 M_\odot$

Stellar Black hole

Wood et al. 2024

# OBSERVATIONS



---

with XMM-Newton

| Multi- $\lambda$ Epochs | X-Shooter VIS, (559.5-1024 nm) | XMM – Newton ~ 0.15 - 12 keV               |
|-------------------------|--------------------------------|--------------------------------------------|
| 1                       | 2023-08-27T00:10:18.635        | -                                          |
| 2                       | 2023-09-04T01:12:32.849        | -                                          |
| 3                       | 2023-09-14T00:41:44.378        | -                                          |
| 4                       | NTT Data 2023-09               | -                                          |
| 5                       | 2023-09-26T00:12:08.807        | -                                          |
| 6                       | 2023-10-01T23:26:04.648        | -                                          |
| 7                       | 2023-10-07T23:48:45.286        | -                                          |
| 8                       | 2023-10-14T00:17:43.073        | -                                          |
| 9                       | 2023-10-17T00:10:29.920        | -                                          |
| 10a                     | 2023-10-21T23:56:36.653        | -                                          |
| 10b                     | 2023-10-22T00:12:49.093        | -                                          |
| 11                      | 2024-02-29T08:54:11.836        | 2024-02-26T14:03:35                        |
| 12                      | 2024-03-24T08:29:31.612        | 2024-03-24T23:26:04<br>2024-03-27T11:57:40 |

See Castro Segura et al.  
2026

# Observations with XMM-Newton



- ★ 3 observations with XMM: 61ks, 20ks, 50ks
- ★ soft-to-hard transition
- ★  $F_x \sim 1.5 \times 10^{-9} \text{ erg s}^{-1} \text{ cm}^{-2}$

# EPIC-PN MODELLING


---

of the EPIC-pn spectra to fit the continuum

# Epoch XMM1

TbFeO \* nthComp \* KYNBB \* xillverCp

Brigitte et al. submitted




$\chi^2 / \text{dof} \approx 1.06$

| Component        | Parameter [units]                 | Value                                       |
|------------------|-----------------------------------|---------------------------------------------|
| <b>TbFeO</b>     | nH [ $10^{22}$ cm <sup>-2</sup> ] | 0.2 (f)                                     |
|                  | O                                 | 0.8 (f)                                     |
|                  | Fe                                | 0.8 (f)                                     |
|                  |                                   |                                             |
| <b>nthComp</b>   | $\Gamma$                          | <b><math>2.1 \pm 0.2</math></b>             |
|                  | kTe [keV]                         | 100 (f)                                     |
|                  | kTbb [keV]                        | 0.3 (f)                                     |
|                  | N                                 | <b><math>0.0186 \pm 5\text{e-}04</math></b> |
| <b>KYNBB</b>     | a/M                               | 0.87 (f)                                    |
|                  | $\theta$ [°]                      | 40.7 ° (f)                                  |
|                  | Rin [rg]                          | <b><math>6.00 \pm 0.05</math></b>           |
|                  | MBH [ $M_{\odot}$ ]               | 10 (f)                                      |
|                  | acc rate [Medd]                   | <b><math>0.012 \pm 0.001</math></b>         |
| <b>xillverCp</b> | log $\xi$                         | —                                           |
|                  | N                                 | —                                           |

# Epoch XXM2

TbFeO \* nthComp \* KYNBB \* xillverCp

Brigitte et al. submitted




$\chi^2 / \text{dof} \approx 1.91$

| Component        | Parameter [units]           | Value                               |
|------------------|-----------------------------|-------------------------------------|
| <b>TbFeO</b>     | nH [ $10^{22}$ cm $^{-2}$ ] | 0.2 (f)                             |
|                  | O                           | 0.7 (f)                             |
|                  | Fe                          | 0.5 (f)                             |
| <b>nthComp</b>   | $\Gamma$                    | <b><math>1.998 \pm 0.006</math></b> |
|                  | kTe [keV]                   | 250 (f)                             |
|                  | kTbb [keV]                  | 0.1 (f)                             |
|                  | N                           | <b><math>0.465 \pm 0.005</math></b> |
| <b>KYNBB</b>     | a/M                         | 0.87 (f)                            |
|                  | $\theta$ [°]                | 40.7 ° (f)                          |
|                  | Rin [rg]                    | <b><math>8.8 \pm 0.3</math></b>     |
|                  | MBH [ $M_{\odot}$ ]         | 10 (f)                              |
|                  | acc rate [Medd]             | <b><math>0.004 \pm 0.001</math></b> |
| <b>xillverCp</b> | log $\xi$                   | <b><math>2.6 \pm 0.4</math></b>     |
|                  | N                           | <b><math>2e-04 \pm 8e-05</math></b> |

# Epoch XMM3


TbFeO \* nthComp \* KYNBB \* xillverCp

Brigitte et al. submitted




$\chi^2 / \text{dof} \approx 1.37$


| Component        | Parameter [units]                | Value                                      |
|------------------|----------------------------------|--------------------------------------------|
| <b>TbFeO</b>     | nH [ $10^{22} \text{ cm}^{-2}$ ] | 0.2 (f)                                    |
|                  | O                                | 0.9 (f)                                    |
|                  | Fe                               | 0.5 (f)                                    |
|                  |                                  |                                            |
| <b>nthComp</b>   | $\Gamma$                         | <b><math>1.74 \pm 0.01</math></b>          |
|                  | kTe [keV]                        | 250 (f)                                    |
|                  | kTbb [keV]                       | 0.1 (f)                                    |
|                  | N                                | <b><math>0.25 \pm 0.02</math></b>          |
| <b>KYNBB</b>     | a/M                              | 0.87 (f)                                   |
|                  | $\theta$ [°]                     | 40.7 ° (f)                                 |
|                  | Rin [rg]                         | <b><math>8.9 \pm 0.1</math></b>            |
|                  | MBH [ $M_{\odot}$ ]              | 10 (f)                                     |
|                  | acc rate [Medd]                  | <b><math>0.004 \pm 0.002</math></b>        |
| <b>xillverCp</b> | log $\xi$                        | <b><math>3.3 \pm 0.1</math></b>            |
|                  | N                                | <b><math>0.001 \pm 1\text{e-}04</math></b> |



$\Gamma \sim 2.1$   
 $N_{\text{Comp}} \sim 0.02$   
 $R_{\text{in}} \sim 6 \text{ rg}$   
 $\text{Acc rate} \sim 0.01$



$\Gamma \sim 1.99$   
 $N_{\text{Comp}} \sim 0.46$   
 $R_{\text{in}} \sim 8.8 \text{ rg}$   
 $\text{Acc rate} \sim 0.004$



$\Gamma \sim 1.74$   
 $N_{\text{Comp}} \sim 0.25$   
 $R_{\text{in}} \sim 8.9 \text{ rg}$   
 $\text{Acc rate} \sim 0.004$

From soft to hard:  $\Gamma$  decreases,  $R_{\text{in}}$  increases and accretion rate decreases

➡ Higher contribution from the corona emission, disk more truncated


# RGS MODELLING

---

of the RGS spectra to fit the absorption and emission lines

Epoch XMM1

TbFeO \* nthComp \* KYNBB \* xillverCp




$\chi^2 / \text{dof} \approx 1.38$

| Component        | Parameter [units]               | Value                 |
|------------------|---------------------------------|-----------------------|
| <b>TbFeO</b>     | nH [ $10^{22} \text{cm}^{-2}$ ] | $0.2 \pm 0.1$         |
|                  | O                               | $0.8 \pm 0.1$         |
|                  | Fe                              | $0.8 \pm 0.1$         |
| <b>nthComp</b>   | $\Gamma$                        | $2.1 \pm 0.2$ (f)     |
|                  | kTe [keV]                       | 100 (f)               |
|                  | kTbb [keV]                      | 0.3 (f)               |
|                  | N                               | 0.0186 (f)            |
| <b>KYNBB</b>     | a/M                             | 0.87 (f)              |
|                  | $\theta$ [°]                    | $40.7^\circ$ (f)      |
|                  | Rin [rg]                        | $6.00 \pm 0.05$ (f)   |
|                  | MBH [ $M_\odot$ ]               | 10 (f)                |
|                  | acc rate [Medd]                 | $0.012 \pm 0.001$ (f) |
| <b>xillverCp</b> | log $\xi$                       | —                     |
|                  | N                               | —                     |

# Epoch XMM2


TbFeO \* nthComp \* KYNBB \* xillverCp




| $\chi^2/\text{dof} \approx 2.0$ |                                  |               |
|---------------------------------|----------------------------------|---------------|
| Component                       | Parameter [units]                | Value         |
| <b>TbFeO</b>                    | nH [ $10^{22} \text{ cm}^{-2}$ ] | $0.2 \pm 0.1$ |
|                                 | O                                | $0.7 \pm 0.1$ |
|                                 | Fe                               | $0.5 \pm 0.2$ |
| <b>nthComp</b>                  | $\Gamma$                         | 1.998 (f)     |
|                                 | kTe [keV]                        | 250 (f)       |
|                                 | kTbb [keV]                       | 0.1 (f)       |
|                                 | N                                | 0.465 (f)     |
| <b>KYNBB</b>                    | a/M                              | 0.87 (f)      |
|                                 | $\theta$ [°]                     | 40.7 ° (f)    |
|                                 | Rin [rg]                         | 8.8 (f)       |
|                                 | MBH [ $M_{\odot}$ ]              | 10 (f)        |
|                                 | acc rate [Medd]                  | 0.004 (f)     |
| <b>xillverCp</b>                | log $\xi$                        | 2.6 (f)       |
|                                 | N                                | 2e-04 (f)     |

# Epoch XMM3


TbFeO \* nthComp \* KYNBB \* xillverCp




| $\chi^2 / \text{dof} \approx 1.15$ |                                  |               |
|------------------------------------|----------------------------------|---------------|
| Component                          | Parameter [units]                | Value         |
| <b>TbFeO</b>                       | nH [ $10^{22} \text{ cm}^{-2}$ ] | $0.2 \pm 0.1$ |
|                                    | O                                | $0.9 \pm 0.1$ |
|                                    | Fe                               | $0.5 \pm 0.2$ |
| <b>nthComp</b>                     | $\Gamma$                         | 1.74 (f)      |
|                                    | kTe [keV]                        | 250 (f)       |
|                                    | kTbb [keV]                       | 0.1 (f)       |
|                                    | N                                | 0.25 (f)      |
| <b>KYNBB</b>                       | a/M                              | 0.87 (f)      |
|                                    | $\theta$ [°]                     | 40.7 ° (f)    |
|                                    | Rin [rg]                         | 8.9 (f)       |
|                                    | MBH [ $M_{\odot}$ ]              | 10 (f)        |
|                                    | acc rate [Medd]                  | 0.004 (f)     |
| <b>xillverCp</b>                   | log $\xi$                        | 3.3 (f)       |
|                                    | N                                | 0.001 (f)     |



$nH \sim 0.26 \times 10^{22} \text{ cm}^{-2}$   
 $O \sim 0.8$   
 $Fe \sim 0.8$



$nH \sim 0.26 \times 10^{22} \text{ cm}^{-2}$   
 $O \sim 0.7$   
 $Fe \sim 0.5$




$nH \sim 0.26 \times 10^{22} \text{ cm}^{-2}$   
 $O \sim 0.9$   
 $Fe \sim 0.5$



ISM absorption consistent except for the Fe abundance,  
most probably instrumental

# RGS SPECTROSCOPY



❖ Absorption lines:

❖ C He

❖ N VI

❖ O edge



❖ Fe I

❖  $V_{\text{shift}} < 100$  km/s

❖ ISM gas and low-ionization gas

Brigitte et al. submitted

RGS



see Castro Segura  
et al. 2026

ption lines:

e

< 100 km/s

as and  
nization gas


mitted

# NEW ATHENA

---

prospects with the X-IFU spectrograph

# SIMULATIONS OF SWIFT J1727 WITH X-IFU




- In the soft-to-hard transition
- 5ks of exposure time
- $F_x \sim 5e-10$  erg/cm<sup>2</sup>/s (2-10 keV)
- SNR  $\sim 42$






- Neon, Magnesium complex
- Emission lines can trace an ionised outflow

0.85–1 keV



## REGIONS OF INTEREST







1.5–2.3 keV

- Silicon complex and Aluminium complex
- O, Ne, Mg, Si absorption edges trace the column density and ionization of the absorbing material

## REGIONS OF INTEREST





## ISM ABSORPTION

- Real variation or from calibration?
- O, Fe abundances
- O edge
- Dust
- Clean separation from intrinsic absorption



## ISM ABSORPTION

- Real variation or from calibration?
- O, Fe abundances
- O edge
- Dust
- Clean separation from intrinsic absorption



## $R_{\text{in}}$ EVOLUTION

- Model dependent
- Truncation evolution in the lower branch of the HID even in transitional states



## ISM ABSORPTION

- Real variation or from calibration?
- O, Fe abundances
- O edge
- Dust
- Clean separation from intrinsic absorption



## $R_{\text{in}}$ EVOLUTION


- Model dependent
- Truncation evolution in the lower branch of the HID even in transitional states



## DISK WIND LINES

- Better spectral resolution for weak winds
- Lower exposure time can access weak lines from weakly-ionised gas

# NEW ATHENA UPCOMING CONFERENCE IN PRAGUE



PRAGUE

NewATHENA science  
conference

2027, April, 19-23  
Institute of Physics of  
Prague



**THANK YOU FOR YOUR ATTENTION**



Image credit: ESA